



## Chesses – fat content

Cheeses are here divided into three classes by fat content.

**Low-fat cheeses**, which can be eaten freely – no limit, are Cottage cheese, Quark, Reduced fat cottage cheese, Reduced fat cheese spread, Reduced fat Edam, Ricotta.

**Medium-fat cheeses** – up to 180g (6oz) per week (if no high fat cheese) are Brie, Camembert, Cheese spread, Danish blue, Edam, Emmental, Feta, Goats milk soft cheese, Medium fat soft cheese, Mozzarella, processed cheese, Reduced fat Cheddar, Reduced fat Cheshire, Smoked processed cheese, Soya cheese.

**High-fat cheeses**, which should be limited to 120g (4oz) per week (if no medium fat cheese), are Caerphilly, Cheddar, Cheshire, Cream cheese, Derby, Double Gloucester, Full fat soft cheese, Gouda, Gruyere, Lancashire, Leicester, Lymeswold, Mascarpone, Parmesan, Red Windsor, Roquefort, Stilton, Vegetarian cheese, Wensleydale.

Medium and high fat cheeses should be eaten only in moderation.

*Either up to 180g (6oz) per week of the medium-fat cheeses,  
or up to 120g (4oz) per week of the high-fat cheeses.*

First published in this form 2002, and updated 2005, 2007, 2008.

All rights reserved. No part of this work may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, without written permission from the BCPA Head Office.

We give permission for copies to be stored and made within the BCPA and any UK hospital; and these hospitals may give printed but not electronic copies to patients provided the source and copyright is acknowledged on the copies – eg include the page footer.